## (\*) Substitution

From the chain rule (for the differentiation of composite functions) we know that

$$\frac{d}{dx}(F(g(x)) = F'(g(x))g'(x).$$

Writing F'(x) = f(x), this gives the corresponding rule for integration

$$\int f(g(x))g'(x) dx = F(g(x)) + C.$$

The difficulty lies in *recognizing* that an integrand is the derivative of a composite function. What we do is look for the following pattern: an integrand that is the product of a composite function and a simpler function, i.e., an integral that looks like this

$$\int f(g(x))k(x)\,dx.$$

If k(x) = g'(x), then we're in business. To compute  $\int f(g(x))g'(x) dx$ , we substitute (rename) u = g(x), which means that du = g'(x) dx so that

$$\int f(g(x))g'(x) dx = \int f(\overbrace{g(x)}^{u}) \overbrace{g'(x)dx}^{du} = \int f(u) du.$$

We still have some work to do, but  $\int f(u)du$  is a simpler integral than the one we started with.

**Example.** To compute  $\int 2x\sqrt{x^2+3}\,dx$ , we first notice that  $(x^2+3)'=2x$ , so that substituting  $u=x^2+3$  means that  $du=2x\,dx$ . This allows us to simplify (and calculate) the integral:

$$\int 2x\sqrt{x^2+3}\,dx = \int \underbrace{\sqrt{x^2+3}}_{\sqrt{2}}\underbrace{2x\,dx}_{2} = \int u^{1/2}\,du = \frac{u^{3/2}}{3/2} + C = \frac{2}{3}(x^2+3)^{3/2} + C.$$

The last step is to replace u by  $x^2 + 3$  in the answer.

## (\*) Adjusting the constant factor

More generally, if in the integral  $\int f(g(x))k(x) dx$ ,  $k(x) = a \cdot g'(x)$  for some (nonzero) constant a, then substituting u = g(x) will still work. In this case we have du = g'(x) dx and since k(x) = ag'(x), it follows that k(x) dx = ag'(x) dx = a(g'(x) dx) = a du, so

$$\int f(g(x))k(x) dx = \int f(g(x)) \overbrace{ag'(x) dx}^{a du} = \int f(u) a du = a \int f(u) du.$$

**Example.** To compute  $\int 5x\sqrt{x^2+3}\,dx$ , we first notice that  $(x^2+3)'=2x$ , so that substituting  $u=x^2+3$  means that  $du=2x\,dx$ . Dividing by 2 shows that  $x\,dx=\frac{1}{2}\,du$  and multiplying by 5 results in the identity  $5x\,dx=\frac{5}{2}\,du$ . Now we can integrate:

$$\int 5x\sqrt{x^2+3}\,dx = \int \underbrace{\sqrt{x^2+3}}_{\sqrt{u}} \underbrace{5x\,dx}_{5x\,dx} = \frac{5}{2} \int u^{1/2}\,du = \frac{5}{2} \cdot \frac{u^{3/2}}{3/2} + C = \frac{5}{3}(x^2+3)^{3/2} + C.$$